COMPATIBILITY OF TWO ADDITIONAL DRUG COMBINATIONS WITH OXYCODONE IN A



SYRINGE PUMP



Colchester Hospital University NHS

IF MORE THAN THREE MEDICATIONS ARE REQUIRED IN THE SYRINGE PUMP PLEASE SEEK SPECIALIST ADVICE

| Compatible | Compatible buse with caution | | | Incompatible | | Avoid – drugs have similar effects | | | Avoid – same class of drug | | Avoid, Antagonis |
|--|------------------------------|-----------|----------------|---------------|--------------------------|------------------------------------|-----------------|----------------|-------------------------------|---------------|---------------------|
| NaCl = Sodium chlor 0.9% WFI = Water for injec | lonaze | Cyclizine | Glycopyrronium | Haloperidol | Hyoscine Butylbromide | Hyoscine hydrobromide | Levomepromazine | Metoclopramide | Midazolam | Octreotide | Ondansetron |
| Clonazepam ^a | | | NaCl | NaCl / WFI | NaCl / WFI | NaCl / WFI | NaCl / WFI | NaCl / WFI | | | |
| Cyclizine | | | | WFIb | | | | | | | |
| Glycopyrronium | NaCl | | | | | | | | | | |
| Haloperidol | NaCl / WFI | WFIb | | | NaCl / WFl | NaCl / WFI | | | NaCl / WFI | NaCl | |
| Hyoscine Butylbrom | NaCL/ | | | NaCl / WFI | | | NaCl / WFI | | NaCl / WFl | | |
| Hyoscine Hydrobrom | NaCL/ | | | NaCl / WFI | | | NaCl / WFI | | NaCl / WFI | | |
| Levomepromazine | NaCl / | | | | NaCl / WFl | NaCl / WFl | | | NaCl / WFI | NaCl / WFI | |
| Metoclopramide | NaCl / WFI | | | | | | | | NaCl / WFI | | С |
| Midazolam | ,,,, | | | NaCl / WFI | NaCl / WFI | NaCl / WFI | NaCl / WFI | NaCl / WFl | | WFI | |
| Octreotide | | | | NaCl | VVI 1 | | NaCl / WFI | *** | WFI | | NaCl / WFI |
| Ondansetron | | | | | | | VVII | С | | NaCl / WFl | VVII |

a = Non PVC tubing must be used with all syringe pumps containing clonazepam

References: Watson M, et al. Oxford Handbook of Palliative Care, 2nd edition, Wyeth. Charlesworth S, et al. Syringe Driver Compatibility Charts.

NAPP pharmaceuticals. Dickman A, et al. The Syringe Driver 2nd edition.

Approved by: Review by: Version 2 Author: Debbie Whittle

b = Crystallisation can occur at higher concentrations, monitor syringe pump regularly.

c = The concurrent use of metoclopramide & ondansetron can increase the risk of cardiac arrhythmias occurring